





ISO9001 & ISO13485 & ISO11137

GelNestTM Matrix, for Organoid Culture, Phenol Red-Free, LDEV-Free

Product overview

GelNestTM Matrix is prepared from basement membrane components extracted from mouse tumor tissues. The main components include laminin, type IV collagen, heparan sulfate proteoglycan, etc. These components can provide the support and signals required for cell adhesion, differentiation, and proliferation. They can also simulate the characteristics of the basement membrane in a physiological environment and improve the success rate and effect of cell culture.

In addition to basement membrane components, GelNestTM Matrix is also rich in a variety of growth factors. These growth factors can promote cell differentiation, proliferation, and migration, further mimicking cell signaling pathways and interactions in physiological environments. GelNestTM Matrix has a wide range of application prospects, especially in tissue engineering, cell culture and research. It can be used for research on organoid culture, stem cell differentiation, angiogenesis, migration or invasion, and *in vivo* tumorigenesis.

Product information

Product number	Product name	Packaging specifications
211282	GelNest™ Matrix, for Organoid Culture,	Bag Package, 5 mL/bottle, 1
	Phenol Red-Free, LDEV-Free	bottle/bag

Product parameters

Source	Mouse tumor tissue basement membrane components
Formulation*	Without phenol red
Protein concentration	Please refer to the COA/COC for batch-specific protein concentration.







ISO9001 & ISO13485 & ISO11137

Appearance	GelNest™ Matrix is liquid at 4°C but forms a gel at 37°C.
Applications	Validated with organoid culture experiment. Suitable for organoid development, cultivation and differentiation.
Storage and shelf life	It is recommended to aliquot the melted product into single-use portions and store it in a -80°C freezer. The product has a shelf life of 2 years.
Precautions	GelNest™ Matrix will start to solidify when the temperature is higher than 10°C. Please operate on ice.

Experimental procedures

Please determine the specific experimental steps based on cell types, culture conditions, and application experience.

Organoid culture

- 1. Re-suspend the single cell suspension used for organoid culture in pre-cooled basal medium at 4°C, and count the cells.
- 2. Mix the cells with GelNestTM Matrix solution and add the mixture to a preheated 24-well plate, each well containing approximately 5x10⁴ cells and 60μL of the matrix gel.
- 3. Immediately place the well plate into the incubator. After about 10 minutes, the matrix gel will solidify.
- 4. Add 500μL of organoid culture medium for culture.
- **5.** Wait 3-5 days for the organoids to form. Finally, the sensitivity of organoids to various drugs can be determined by imaging live cells through high-content microscopy.





ISO9001 & ISO13485 & ISO11137




Figure 1. Growth of human cholangiocarcinoma organoids in Brand C (top) and GelNestTM matrix gel (bottom) for 6 days. Scale bar represents 200μm.

Safety recommendations and limitations

Please follow good laboratory safety practices.

For research use only. Not intended for diagnostic or therapeutic purposes. Contains ingredients of animal origin.

Technical support and contact information

For FAQ, GelNest™ Matrix Selection Guide, Quality Assurance COA/COC or other technical support and product issues, please refer to our website or use the following contact information.

Production and after-sales service unit: Wuxi NEST Biotechnology Co., Ltd.

Production and after-sales address: No.530, Xida Road, Meicun Industrial Park, Xinwu District, Wuxi, Jiangsu, China

Tel: +86-510-68006788

Email: info@nest-wuxi.com

Website: www.cell-nest.com